

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

\$%*STN:HighlightOn= ***;HighlightOff=***;
Trying 3106016892...Open

Welcome to STN International! Enter xx
LOGINID:asspta1633xq
PASSWORD:
TERMINAL (ENTER 1, 2, 3, OR 7):2

***** Welcome to STN International *****

NEWS 1 Web Page URLs for STN Seminar Schedule - N. America
NEWS 2 Dec 17 The CA Lexicon available in the CAPLUS and CA files
NEWS 3 Feb 06 Engineering Information Encompass files have new names
NEWS 4 Feb 16 TOXLINE no longer being updated
NEWS 5 Apr 23 Search Derwent WPIINDEX by chemical structure
NEWS 6 Apr 23 PRE-1967 REFERENCES NOW SEARCHABLE IN CAPLUS AND CA
NEWS 7 May 07 DGENE Reload
NEWS 8 Jun 20 Published patent applications (A1) are now in USPATFULL
NEWS 9 JUL 13 New SDI alert frequency now available in Derwent's
DWPI and DPCI
NEWS 10 Aug 23 In-process records and more frequent updates now in
MEDLINE
NEWS 11 Aug 23 PAGE IMAGES FOR 1947-1966 RECORDS IN CAPLUS AND CA
NEWS 12 Aug 23 Ads Newsletters (ADISNEWS) now available on STN
NEWS 13 Sep 17 IMSworld Pharmaceutical Company Directory name change
to PHARMASEARCH
NEWS 14 Oct 08 Korean abstracts now included in Derwent World Patents
Index
NEWS 15 Oct 08 Number of Derwent World Patents Index updates increased
NEWS 16 Oct 15 Calculated properties now in the REGISTRY/REGISTRY File
NEWS 17 Oct 22 Over 1 million reactions added to CASREACT
NEWS 18 Oct 22 DGENE GETSIM has been improved
NEWS 19 Oct 29 AAASD no longer available
NEWS 20 Nov 19 New Search Capabilities USPATFULL and USPAT2
NEWS 21 Nov 19 TOXCENTER(SM) - new toxicology file now available on STN
NEWS 22 Nov 29 COPPERLIT now available on STN
NEWS 23 Nov 29 DWPI revisions to NTIS and US Provisional Numbers
NEWS 24 Nov 30 Files VETU and VETB to have open access
NEWS 25 Dec 10 WPIINDEXAWPIDSAWPIX New and Revised Manual Codes for 2002
NEWS 26 Dec 10 DGENE BLAST Homology Search
NEWS 27 Dec 17 WELDASEARCH now available on STN
NEWS 28 Dec 17 STANDARDS now available on STN
NEWS 29 Dec 17 New fields for DPCI
NEWS 30 Dec 19 CAS Roles modified
NEWS 31 Dec 19 1907-1948 data and page images added to CA and Caphus

NEWS EXPRESS August 15 CURRENT WINDOWS VERSION IS V8.0c,
CURRENT MACINTOSH VERSION IS V8.0 (ENG) AND V6.0J (JP),
AND CURRENT DISCOVER FILE IS DATED 07 AUGUST 2001

NEWS HOURS STN Operating Hours Plus Help Desk Availability
NEWS INTER General Internet Information
NEWS LOGIN Welcome Banner and News Items
NEWS PHONE Direct Dial and Telecommunication Network Access to STN
NEWS WWW CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that
specific topic.

All use of STN is subject to the provisions of the STN Customer
agreement. Please note that this agreement limits use to scientific
research. Use for software development or design or implementation
of commercial gateways or other similar uses is prohibited and may
result in loss of user privileges and other penalties.

*****STN Columbus*****

FILE 'HOME' ENTERED AT 16:29:22 ON 02 JAN 2002

=> file biosis embase caplus
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 0.15 0.15

FILE 'BIOSIS' ENTERED AT 16:29:46 ON 02 JAN 2002
COPYRIGHT (C) 2002 BIOSIS(R)

FILE 'EMBASE' ENTERED AT 16:29:48 ON 02 JAN 2002
COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'CAPLUS' ENTERED AT 16:29:48 ON 02 JAN 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE 'HELP USAGETERMS' FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> s adeno-associated viruses or AAV
L1 3153 ADENO-ASSOCIATED VIRUSES OR AAV

=> s adeno associated virus? or AAV
L2 3153 ADENO ASSOCIATED VIRUSE? OR AAV

=> s adeno associated virus 2 or AAV-2
L3 536 ADENO ASSOCIATED VIRUS 2 OR AAV-2

=> s l1 or l3

L4 3233 L1 OR L3

=> s l4 and cancer?

L5 201 L4 AND CANCER?

=> s l4 and chemo? resistan?

L6 5 L4 AND CHEMO? RESISTANT?

=> dup rem l6

PROCESSING COMPLETED FOR L6

L7 2 DUP REM L6 (3 DUPLICATES REMOVED)

=> d bib abs l-

L- IS NOT A VALID FORMAT FOR FILE 'BIOSIS'

The following are valid formats:

The default display format is BIB.

ABS ----- AB
ALL ----- AN, DN, TI, AU, CS, PI, SO, NTE, DT, FS, LA, SL,
AB, NCL, CC, BC, IT (CT, ST), CO, NA, GT, ORGN, RN
BIB ----- AN, DN, TI, AU, CS, PI, SO, NTE, DT, FS, LA, SL
CBIB ----- AN, compressed bibliographic information
DALL ----- ALL, delimited for post-processing
IABS ----- ABS, with text label
IALL ----- ALL, indented with text labels
IBIB ----- BIB, indented with text labels
IIND ----- IND, indented with text labels
IND ----- NCL, CC, BC, IT (CT, ST), CO, NA, NP, GT, ORGN, RN

HIT ----- all fields containing hit search terms
HITIND ----- IND
KWIC ----- Hit terms plus 20 words on either side
OCC ----- number of occurrences of hit terms and fields
in which they occur

Hit terms will be highlighted in all available fields except MY and
PY.

To display a particular field or fields, enter the display field
codes. For a list of display field codes, enter 'HELP DFIELDS'
at an arrow prompt (=>). Examples of formats include: 'BIB',
'AB', 'SO, ST'. You may specify the format fields in any order,
and the information will be displayed in the same order as the
format specification.

The same formats (except for HIT, HITIND, KWIC, and OCC) may be
used with the DISPLAY ACC command to display the record for a
specified Accession Number.
ENTER DISPLAY FORMAT (BIB):bib

L7 ANSWER 1 OF 2 BIOSIS COPYRIGHT 2002 BIOSIS DUPLICATE 1
AN 1999:192567 BIOSIS
DN PREV199900192567
TI Enhanced sensitivity of small cell lung cancer cell lines to cisplatin and
etoposide after infection with adeno-associated virus type 2.
AU Hilgenberg, M.; Schlehofer, J. R.; von Knebel Doeberitz, M.;
Klein-Baumenschmidt, P. (1)
CS (1) Sektion fuer Molekulare Diagnostik und Therapie, Chirurgische
Universitaetsklinik, Universitaet Heidelberg, INF 120, 69120, Heidelberg
Germany
SO European Journal of Cancer, (Jan., 1999) Vol. 35, No. 1, pp. 108-110.
ISSN: 0959-8049.
DT Article
LA English

=> d bib abs l-
YOU HAVE REQUESTED DATA FROM 2 ANSWERS - CONTINUE? Y(N):y

L7 ANSWER 1 OF 2 BIOSIS COPYRIGHT 2002 BIOSIS DUPLICATE 1
AN 1999:192567 BIOSIS
DN PREV199900192567
TI Enhanced sensitivity of small cell lung cancer cell lines to cisplatin and
etoposide after infection with adeno-associated virus type 2.
AU Hilgenberg, M.; Schlehofer, J. R.; von Knebel Doeberitz, M.;
Klein-Baumenschmidt, P. (1)
CS (1) Sektion fuer Molekulare Diagnostik und Therapie, Chirurgische
Universitaetsklinik, Universitaet Heidelberg, INF 120, 69120, Heidelberg
Germany
SO European Journal of Cancer, (Jan., 1999) Vol. 35, No. 1, pp. 108-110.
ISSN: 0959-8049.
DT Article
LA English
AB In previous studies we have reported the sensitisation of human tumour
cells to gamma irradiation and chemotherapeutic drugs upon infection with
the human non-pathogenic adeno-associated virus type 2 ("AAV" -
"2") in vitro and in vivo. Treatment of small cell lung cancer
(SCLC) is consistently hampered by relapses due to the selection of
"chemotherapy" - "resistant" cell clones. Hence, we were
interested to test whether selection of "chemotherapy" -
"resistant" SCLC cells might be reduced or even prevented if
chemotherapy is applied in combination with "AAV" - "2"
infection. In vitro proliferation assays indicated that the number of
proliferating cells, after combined treatment with cisplatin and
etoposide, can be significantly reduced by concomitant "AAV" -
"2" infection, as compared with treated but non-infected controls.
H460 SCLC cells, which show resistance to etoposide/cisplatin chemotherapy
(compared with a cell line which was never chemotherapeutically treated
before, like NCH-H209) were significantly more sensitive after "AAV"
"2" infection, suggesting that the therapeutic efficacy of
chemotherapy in SCLC can be enhanced even if the cells are already
relatively resistant to chemotherapy. Similarly, in vivo growth of tumours
induced by inoculation of SCLC cells into immunocompromised nude mice was
reduced more efficiently in "AAV" - "2" -infected animals
compared with tumours in mice treated with chemotherapeutic drugs alone.
These data extend and further support our previous reports on "AAV"
functions which might be useful in improving the efficacy of
chemotherapeutic drugs used in human cancer treatment.

L7 ANSWER 2 OF 2 BIOSIS COPYRIGHT 2002 BIOSIS DUPLICATE 2
AN 1997:87547 BIOSIS
DN PREV199789378260
TI Gene therapy for haematopoietic and lymphoid disorders.
AU Kohn, D. B.
CS Div. Res. Immunol. Bone Marrow Transplantation, Child. Hosp. Los Angeles,
Mailstop 62, 4650 Sunset Boulevard, Los Angeles, CA 90027 USA
SO Clinical and Experimental Immunology, (1997) Vol. 107, No. SUPPL. 1, pp.
54-57
ISSN: 0009-9104.
DT Article
LA English
AB Gene transfer into haematopoietic stem cells (HSC) has been investigated
for treatment of genetic disorders, conferral of "chemotherapy"
"resistance" and insertion of genes to inhibit HIV-1 replication.

Methods have been available for almost a decade to transduce murine HSC using high-titre retroviral vectors and stimulation of HSC proliferation with cytokines such as IL-3 and IL-6. Unfortunately, attempts to replicate the high efficiency of gene transfer using canine or simian gene transferbone marrow transplantation models have consistently shown that only a small fraction (0.1-1%) of reconstituting HSC are transduced using protocols similar to those which are successful in murine models. Initial clinical trials using retroviral-mediated gene transfer into human HSC also produced minimal transduction frequencies. The dichotomous results may reflect differences in the cell cycle kinetics of murine HSC versus those of larger mammals or the density of receptors for the retroviral vectors on the cells. Attempts to increase the fraction of HSC which are in active cell cycle, a prerequisite for retroviral-mediated transduction, have used either combinations of recombinant cytokines, culture on marrow stromal layers, or alternative sources for HSC; such as mobilized peripheral blood stem cells or umbilical cord blood. Other efforts have used retroviral vectors packaged with either the Gibbon Ape Leukemia virus envelope or the Vesicular Stomatitis Virus G protein. To date, none of these methods has produced a significantly increased frequency of long-term reconstituting HSC. Results using adeno-associated virus (AAV)-based vectors for HSC transduction have been conflicting, with the stable persistence of non-integrated virus particles making interpretation of results difficult using in vitro assays. Therefore, clinical trials may best be directed toward disorders that may benefit from a small fraction of genetically corrected HSC. These would include disorders where progeny of corrected HSC would be expected to have a selective survival advantage (e.g. SCID, WAS, HIV, ? chemoresistance) or where a small fraction of corrected cells can have a direct clinical benefit (e.g. CGD, MPS). Further basic research into HSC biology and gene delivery vectors must continue for wider application, such as haemoglobinopathies and some lysosomal storage diseases.

=> FIL STNGUIDE
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 41.57 41.72

FILE 'STNGUIDE' ENTERED AT 16:36:04 ON 02 JAN 2002
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: Dec 14, 2001 (20011214UP).

=> FIL EMBASE BIOSIS CAPLUS
COST IN U.S. DOLLARS SINCE FILE TOTAL
ENTRY SESSION
FULL ESTIMATED COST 0.00 41.72

FILE 'EMBASE' ENTERED AT 16:38:10 ON 02 JAN 2002
COPYRIGHT (C) 2002 Elsevier Science B.V. All rights reserved.

FILE 'BIOSIS' ENTERED AT 16:38:10 ON 02 JAN 2002
COPYRIGHT (C) 2002 BIOSIS(R)

FILE 'CAPLUS' ENTERED AT 16:38:10 ON 02 JAN 2002
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
PLEASE SEE 'HELP USAGETERMS' FOR DETAILS.
COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

=> d his

(FILE 'HOME' ENTERED AT 16:29:22 ON 02 JAN 2002)

FILE 'BIOSIS, EMBASE, CAPLUS' ENTERED AT 16:29:46 ON 02 JAN 2002
L1 3153 S ADENO-ASSOCIATED VIRUSES OR AAV
L2 3153 S ADENO ASSOCIATED VIRUSE? OR AAV
L3 538 S ADENO ASSOCIATED VIRUS 2 OR AAV-2
L4 3233 S L1 OR L3
L5 201 S L4 AND CANCER?
L6 5 S L4 AND CHEMO? RESISTAN?
L7 2 DUP REM L6 (3 DUPLICATES REMOVED)

FILE 'STNGUIDE' ENTERED AT 16:36:04 ON 02 JAN 2002

FILE 'EMBASE, BIOSIS, CAPLUS' ENTERED AT 16:38:10 ON 02 JAN 2002

=> s l4 and (colon cancer or pancreatic carcinoma or brain tumor or small cell lung cancer)
L8 10 L4 AND (COLON CANCER OR PANCREATIC CARCINOMA OR BRAIN TUMOR OR SMALL CELL LUNG CANCER)

=> s l4 and radiotherapy and resistance
L9 0 L4 AND RADIOTHERAPY AND RESISTANCE

=> s l4 and resistance
L10 170 L4 AND RESISTANCE

=> s l10 and (chemo or radiot?)
L11 1 L10 AND (CHEMO OR RADIOT?)

=> s l10 and radiation
L12 5 L10 AND RADIATION

=> s l11 or l12
L13 6 L11 OR L12

=> dup rem l13
PROCESSING COMPLETED FOR L13
L14 4 DUP REM L13 (2 DUPLICATES REMOVED)

=> d bib abs 1-
YOU HAVE REQUESTED DATA FROM 4 ANSWERS - CONTINUE? Y(N):y

L14 ANSWER 1 OF 4 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V. DUPLICATE 1
AN 2001287672 EMBASE
TI A hypoxia-regulated adeno-associated virus vector for cancer-specific gene therapy.
AU Ruan H.; Su H.; Hu L.; Lamborn K.R.; Kan Y.W.; Deen D.F.
CS Dr. D.F. Deen, Brain Tumor Research Center, University of California, San Francisco, CA 94143-0520, United States. ddeen@itsa.ucsf.edu

SO Neoplasia, (2001) 3/3 (255-263).
Refs: 35

ISSN: 1522-8002 CODEN: NEOPFL

CY United States

DT Journal; Article

FS 018 Cancer

022 Human Genetics

LA English

SL English

AB The presence of hypoxic cells in human brain tumors is an important factor leading to "resistance" to "radiation" therapy. However, this physiological difference between normal tissues and tumors also provides the potential for designing cancer-specific gene therapy. We compared the increase of gene expression under anoxia (<0.01% oxygen) produced by 3, 6, and 9 copies of hypoxia-responsive elements (HRE) from the erythropoietin gene (Epo), which are activated through the transcriptional complex hypoxia-inducible factor 1 (HIF-1). Under anoxic conditions, nine copies of HRE (9XHRE) yielded 27- to 37-fold of increased gene expression in U-251 MG and U-87 MG human brain tumor cell lines. Under the less hypoxic conditions of 0.3% and 1% oxygen, gene activation by 9XHRE increased expression 11- to 18-fold in these cell lines. To generate a recombinant adeno-associated virus (rAAV) in which the transgene can be regulated by hypoxia, we inserted the DNA fragment containing 9XHRE and the LacZ reporter gene into an "AAV" vector. Under anoxic conditions, this vector produced 79- to 110-fold increase in gene expression. We believe this hypoxia-regulated rAAV vector will provide a useful delivery vehicle for cancer-specific gene therapy.

L14 ANSWER 2 OF 4 BIOSIS COPYRIGHT 2002 BIOSIS

AN 2002264838 BIOSIS

DN PREV200002264838

TI Enhanced sensitivity of pancreatic tumour cells to 5-FU-chemotherapy mediated by adeno-associated virus type 2 ("AAV" - "2") infection in vitro and in vivo.

AU Eisold, Sven Christian (1); Ridder, Ruediger; Ryschisch, Eduard; Schmidt, Jan; Meyer, Geske C.; Klar, Ernst; Herfarth, Christian; von Knebel-Doeberitz, Magnus

CS (1) Dept Surg, Univ of Heidelberg, Heidelberg Germany

SO Gastroenterology, (April, 2000) Vol. 116, No. 4 Suppl. 2 Part 1, pp. AGA

A531, print.

Meeting Info.: 101st Annual Meeting of the American Gastroenterological Association and the Digestive Disease Week, San Diego, California, USA May 21-24, 2000 American Gastroenterological Association

ISSN: 0018-5085.

DT Conference

LA English

SL English

L14 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2002 ACS

AN 1996642183 CAPLUS

DN 125:292439

TI Improved efficacy of chemotherapy by parvovirus-mediated sensitization of human tumor cells

AU Klein-Baumenschmitt, P.; Von Knebel Doeberitz, M.; Ehrbar, M.; Geletheky, K.; Kleinschmidt, J.; Schlehner, J. R.

CS German Cancer Research Center, Applied Tumor Virology, Heidelberg, 69120, Germany

SO Eur. J. Cancer, Part A (1996), 32A(10), 1774-1780

CODEN: EJCTEA

DT Journal

LA English

AB Increasing "resistance" of tumor cells towards the cytotoxic action of chemotherapeutic drugs is a major limitation in the treatment of cancer patients. The non-pathogenic human adeno-assoc. viruses ("AAV") have been reported to sensitize HeLa cervical cancer cells to gamma irradiation in vivo and in vitro. To test whether these parvoviruses might render other human tumor cells more sensitive towards chemotherapeutic drugs, we analyzed the effects of "AAV" type 2 ("AAV" - "2") infection on established cancer cell lines and freshly explanted tumor biopsies treated with chemotherapeutic agents (e.g. cisplatin). "AAV" - "2" infection significantly increased the cytotoxic activity of chemotherapeutic drugs compared with uninfected controls. "AAV" - "2" infection without concomitant chemotherapeutic treatment had no significant effect on viability of the cells. In nude mice, combined application of "AAV" - "2" infection and chemotherapeutic treatment significantly increased the therapeutic activity on tumors arising from s.c. injected tumor cells compared with tumors treated by chemotherapeutics only. These results indicate that "AAV" - "2" infection sensitizes human cancer cells towards the cytotoxic action of chemotherapeutic drugs.

L14 ANSWER 4 OF 4 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.

AN 92302498 EMBASE

DN 1992302498

TI Modulation of the cellular phenotype by integrated adeno-associated virus.

AU Winocour E.; Puzis L.; Eldin S.; Koch T.; Danovitch B.; Mendelson E.; Shaulian E.; Karby S.; Lav S.

CS Dept. of Molecular Genetics/Virology, Weizmann Institute of Science, Rehovot, Israel

SO Virology, (1992) 190(1) (318-329).

ISSN: 0042-6822 CODEN: VIRLAX

CY United States

DT Journal; Article

FS 004 Microbiology

037 Drug Literature Index

LA English

SL English

AB The adeno-associated virus ("AAV") rep gene encodes a series of overlapping, multifunctional, nonstructural proteins (Rep proteins) which regulate the viral life cycle and which are also capable of trans-regulating nonviral gene expressions (reviewed in Berns, 1990, Microbiol. Rev. 54, 318-328). To investigate the expression of the "AAV" rep gene in a cellular chromosomal context, SV40-transformed Chinese hamster embryo (OD4) cells were infected with an "AAV" /neo hybrid virus and progeny resistant to the antibiotic G418 were selected and amplified. Chromosomal integration and RNA transcription of the "AAV" and neo DNA inserts were confirmed by Southern and Northern blotting procedures. One of the G418(R) cell lines stably expressed a protein which reacted specifically with "AAV" anti-Rep antiserum in Western immunoblots. The stable integration of "AAV" rep DNA, which did not interfere with cell proliferation under normal growth conditions, was associated with two changes in cellular phenotype: eight

of nine lines were markedly more sensitive to UV light (254 nm) than were the parental OD4 cells; and seven of the nine lines had lost the capacity to promote SV40 origin DNA amplification in vitro, in contrast to the parental OD4 cells. OD4 cells transformed to G418(R) by ***AAV*** /neo DNA constructs with a deleted rep gene, or by a neo DNA construct lacking ***AAV*** DNA, did not display these phenotypic changes. It is suggested that stable integration of the ***AAV*** rep gene interferes with cellular processes connected with DNA repair and gene amplification.

=> d his

(FILE 'HOME' ENTERED AT 18:29:22 ON 02 JAN 2002)

FILE 'BIOSIS, EMBASE, CAPLUS' ENTERED AT 18:29:46 ON 02 JAN 2002

L1 3153 S ADENO-ASSOCIATED VIRUSES OR AAV
L2 3153 S ADENO ASSOCIATED VIRUS? OR AAV
L3 536 S ADENO ASSOCIATED VIRUS 2 OR AAV-2
L4 3233 S L1 OR L3
L5 201 S L4 AND CANCER?
L6 5 S L4 AND CHEMO? RESISTANT?
L7 2 DUP REM L6 (3 DUPLICATES REMOVED)

FILE 'STNGUIDE' ENTERED AT 18:38:04 ON 02 JAN 2002

FILE 'EMBASE, BIOSIS, CAPLUS' ENTERED AT 18:38:10 ON 02 JAN 2002

L8 16 S L4 AND COLON CANCER OR PANCREATIC CARCINOMA OR BRAIN TUMOR
L9 9 S L4 AND RADIOTHERAPY AND RESISTANCE
L10 170 S L4 AND RESISTANCE
L11 1 S L10 AND (CHEMO OR RADIOT?)
L12 5 S L10 AND RADIATION
L13 8 S L11 OR L12
L14 4 DUP REM L13 (2 DUPLICATES REMOVED)

=> dup rem l8

PROCESSING COMPLETED FOR L8

L15 9 DUP REM L8 (7 DUPLICATES REMOVED)

=> d bib abs 1-

YOU HAVE REQUESTED DATA FROM 9 ANSWERS - CONTINUE? Y(N):y

L15 ANSWER 1 OF 9 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 1

AN 2001287072 EMBASE

TI A hypoxia-regulated adeno-associated virus vector for cancer-specific gene therapy.

AU Ruan H.; Su H.; Hu L.; Lamborn K.R.; Kan Y.W.; Deen D.F.
CS Dr. D.F. Deen, Brain Tumor Research Center, University of California, San Francisco, CA 94143-0520, United States. ddeen@itsa.ucsf.edu
SO Neoplasia, (2001) 3/3 (255-263).

Refs: 35

ISSN: 1522-8002 CODEN: NEOPFL

CY United States

DT Journal Article

FS 016 Cancer

022 Human Genetics

LA English

SL English

AB The presence of hypoxic cells in human brain tumors is an important factor leading to resistance to radiation therapy. However, this physiological difference between normal tissues and tumors also provides the potential for designing cancer-specific gene therapy. We compared the increase of gene expression under anoxia (<0.01% oxygen) produced by 3, 6, and 9 copies of hypoxia-responsive elements (HRE) from the erythropoietin gene (Epo), which are activated through the transcriptional complex hypoxia-inducible factor 1 (HIF-1). Under anoxic conditions, nine copies of HRE (9X-HRE) yielded 27- to 37-fold of increased gene expression in U-251 MG and U-87 MG human ***brain*** ***tumor*** cell lines. Under the less hypoxic conditions of 0.3% and 1% oxygen, gene activation by 9X-HRE increased expression 11- to 18-fold in these cell lines. To generate a recombinant adeno-associated virus (AAV) in which the transgene can be regulated by hypoxia, we inserted the DNA fragment containing 9X-HRE and the LacZ reporter gene into an ***AAV*** vector. Under anoxic conditions, this vector produced 79- to 110-fold increase in gene expression. We believe this hypoxia-regulated rAAV vector will provide a useful delivery vehicle for cancer-specific gene therapy.

L15 ANSWER 2 OF 9 CAPLUS COPYRIGHT 2002 ACS

AN 2000:68258 CAPLUS

DN 133:26253

TI Prolactin regulatory element binding (PREB) protein, cDNA and genetic sequences and therapeutic and diagnostic uses thereof

IN Bancroft, Carter F.; Fless, Makiko; Clelland, Catherine L.

PA Mount Sinai School of Medicine of New York University, USA

SO PCT Int. Appl., 87 pp.

CODEN: PIXX02

DT Patent

LA English

FAN CNT 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI WO 2000/056756	A2	20000928	WO 2000-US7642	20000323

W: CA, JP, US

RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE

PRAI US 1999-125728 P 19990323

AB The present invention relates to the discovery of a novel transcription factor called PREB (prolactin regulatory element binding) protein which functions in the kinase-mediated hormonal regulation of prolactin gene expression. The invention based on the discovery and characterization of the rat Preb gene and protein and the identification of a cloned human DNA contig, the human PREB gene. The invention provides for an assay for distinguishing between different ***brain*** ***tumor*** types whereby PREB transcript levels are quantified. The presence of PREB would be indicative the presence of astrocytoma ***brain*** ***tumor*** cells but not neuroepithelioma or glioma ***brain*** ***tumor*** cells. Also the present invention provides for methods for the treatment of cancers and autoimmune diseases through the inhibition of prolactin gene expression.

L15 ANSWER 3 OF 9 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 2

AN 2000117450 EMBASE

TI Current status of viral gene therapy for brain tumors.

AU Gupta N.
CS N. Gupta, Univ. Chicago Children's Hospital, 5841 South Maryland Ave., Chicago, IL 60637, United States. ngupta@peds.bsd.uchicago.edu
SO Expert Opinion on Investigational Drugs, (2000) 9/4 (713-726).

Refs: 94

ISSN: 1354-3784 CODEN: EOIDER

CY United Kingdom

DT Journal, General Review

FS 008 Neurology and Neurosurgery

018 Cancer

022 Human Genetics

LA English

SL English

AB Malignant glial tumours represent the majority of primary brain tumours. Despite the use of many adjunctive treatment strategies in addition to surgery, the prospect of cure or even long-term survival is poor. In the last decade, there has been an explosion of interest in the development of delivery systems that will allow the expression of exogenous genes in the CNS. For the most part, these systems are based upon modified viruses. To date, the greatest experience has been with retroviruses, herpes simplex virus 1 (HSV), adenovirus and adeno-associated virus (***AAV***). This review will outline the biology of these viral vectors, modifications permitting in vivo administration and their respective advantages and disadvantages for the treatment of malignant brain tumours. The present obstacles to gene therapy strategies will also be described. To date, no convincing clinical trial has emerged that provides objective proof of the superiority of gene therapy strategies as compared to conventional treatment.

L15 ANSWER 4 OF 9 BIOSIS COPYRIGHT 2002 BIOSIS

AN 2001:122449 BIOSIS

DN PREV200100122449

TI Host range of adeno-associated viral (***AAV***) vectors and implications for ***AAV*** tumor gene therapy.

AU Veldwijk M. R. (1); Kleinschmidt, J. A. (1); Zeller, W. J. (1); Fruehauf, S.

CS (1) German Cancer Research Center (DKFZ) Heidelberg, Heidelberg Germany

SO Onkologie, (October, 2000) Vol. 23, No. Sonderheft 7, pp. 176. print.

Meeting Info.: Annual Meeting of the German and Austrian Society for Hematology and Oncology Graz, Austria October 21-25, 2000

ISSN: 0378-584X.

DT Conference

LA English

SL English

L15 ANSWER 5 OF 9 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 3

AN 1999254105 EMBASE

TI Superior gene transfer into solid tumour cells than into human mobilised peripheral blood progenitor cells using helpervirus-free adeno-associated viral vector stocks.

AU Veldwijk M.R.; Schiedmeier B.; Kleinschmidt J.A.; Zeller W.J.; Fruehauf S.

CS M.R. Veldwijk, German Cancer Research Centre (DKFZ), Im Neuenheimer Feld

280, D-69120 Heidelberg, Germany. m.veldwijk@dkfz-heidelberg.de

SO European Journal of Cancer, (1999) 35/7 (1136-1142).

Refs: 29

ISSN: 0959-8049 CODEN: EJCAEL

PUI S 0959-8049(99)00075-1

CY United Kingdom

DT Journal Article

FS 016 Cancer

022 Human Genetics

037 Drug Literature Index

LA English

SL English

AB Autologous peripheral blood progenitor cell (PBPC) grafts can be contaminated with tumour cells that potentially give rise to relapse following myeloablative therapy and PBPC transplantation. Adeno-associated virus (***AAV***)-based vectors produced by a new adenovirus-free technique are a gene delivery system which may be applicable for tumour cell purging. To test for the host range of these vectors, solid tumours of clinical relevance and normal GD34+ PBPG were selected as target cells for an ***AAV***-vector, encoding the green-fluorescent protein (GFP) as the indicator gene. At a multiplicity of infection (MOI) of 100/79.94% +/- 14.36% (mean +/- SEM) of the connective tissue sarcoma cell line (HS-1) and 84.84% +/- 6.91% of the cervical carcinoma cell line (HeLa-RG) expressed GFP while the other cell lines tested (1 ovarian tumour, 1 germ cell tumour, 1 osteosarcoma, 2 ***small*** ***cell*** ***lung*** ***cancer***) ranged between 2.62% and 11.94%. Optimising the transduction protocol by use of higher MOIs of up to 500 and by pretreatment with the tyrosine kinase inhibitor, genistein, resulted in up to 95.97% and 94.10% green-fluorescent HS-1 and HeLa-RG cells, respectively. In contrast, only 1.39% +/- 0.51% of the normal haematopoietic GD34+ progenitor cells expressed GFP at a MOI of 100. The differential infectivity between HS-1 and GD34+ cells was maintained after tumour cell spiking in leucapheresis products. Our observations suggest that ***AAV***-based vectors may prove useful for purging of autologous PBPG grafts from solid tumour cells.

L15 ANSWER 6 OF 9 CAPLUS COPYRIGHT 2002 ACS

AN 2000:24267 CAPLUS

DN 132:288168

TI HSV-1-based vectors for gene therapy of neurological diseases and brain tumors: Part II. Vector systems and applications

AU Jacobs, Andreas; Breakfield, Xandra O.; Fraefel, Cornel

CS Department of Neurology at the University, MPI for Neurological Research, Cologne, 50931, Germany

SO Neoplasia (N. Y.) (1999) 1(5), 402-416

CODEN: NEOPFL; ISSN: 1522-8002

PB Stockton Press

DT Journal, General Review

LA English

AB A review with 202 refs. Many properties of HSV-1 are esp. suitable for using this virus as a vector to treat diseases affecting the central nervous system (CNS), such as Parkinson's disease or malignant gliomas. These advantageous properties include natural neurotropism, high transduction efficiency, large transgene capacity, and the ability of entering a latent state in neurons. Selective oncolysis in combination with modulation of the immune response mediated by replication-conditional HSV-1 vectors appears to be a highly promising approach in the battle against malignant glioma. Helper virus-free HSV/ ***AAV*** hybrid amplicon vectors have great promise in mediating long-term gene expression

in the PNS and CNS for the treatment of various neurodegenerative disorders or chronic pain. Current research focuses on the design of HSV-1-derived vectors which are targeted to certain cell types and support transcriptionally regulatable transgene expression. Here, we review the recent developments on HSV-1-based vector systems and their applications in exptl. and clin. gene therapy protocols.

RE.CNT 202
RE
(2) Aboody-Guterman, K; NeuroReport 1997, V8, P3801 CAPLUS
(3) Adams, P; Semin Cancer Biol 1995, V6, P98 CAPLUS
(4) Adams, P; Semin Cancer Biol 1995, V6, P98 CAPLUS
(5) Agri, M; J Natl Cancer Inst 1998, V90, P370 CAPLUS
(7) Andersen, J; Cell Mol Neurobiol 1993, V13, P503 CAPLUS
ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 7 OF 9 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 4
AN 1999082500 EMBASE
TI Enhanced sensitivity of ***small*** ***cell*** ***lung***
cancer cell lines to cisplatin and etoposide after infection with
adeno-associated virus type 2
AU Hilgenberg M; Schlehofer J.R.; Von Knebel Doeberitz M.; Klein-
Baumenschmitt P.
CS P. Klein-Baumenschmitt, Sekl. Molekulare Diagnostik Therapie, Chirurgische
Universitätsklinik, Universität Heidelberg, 69120 Heidelberg, Germany.
p.klein@kldzheidelberg.de
SO European Journal of Cancer, (1999) 35/1 (106-110).
Refs: 21
ISSN: 0959-8049 CODEN: EJCAEL
PUI: S 0959-8049(98)00275-5
CY United Kingdom
DT Journal; Article
FS 004 Microbiology
015 Chest Diseases, Thoracic Surgery and Tuberculosis
016 Cancer
037 Drug Literature Index
LA English
SL English
AB In previous studies we have reported the sensitisation of human tumour
cells to gamma irradiation and chemotherapeutic drugs upon infection with
the human non-pathogenic adeno-associated virus type 2 (***AAV***-
2). In vitro and in vivo. Treatment of ***small*** ***cell***
lung ***cancer*** (SCLC) is consistently hampered by relapses
due to the selection of chemotherapy-resistant cell clones. Hence, we were
interested to test whether selection of chemotherapy-resistant SCLC cells
might be reduced or even prevented if chemotherapy is applied in
combination with ***AAV***-***2*** infection. In vitro
proliferation assays indicated that the number of proliferating cells,
after combined treatment with cisplatin and etoposide, can be
significantly reduced by concomitant ***AAV***-***2*** infection,
as compared with treated but non-infected controls. H460 SCLC cells, which
show resistance to etoposide/cisplatin chemotherapy (compared with a cell
line which was never chemotherapeutically treated before, like NCI-H209)
were significantly more sensitive after ***AAV***-***2***
infection, suggesting that the therapeutic efficacy of chemotherapy in
SCLC can be enhanced even if the cells are already relatively resistant to
chemotherapy. Similarly, in vivo growth of tumors induced by inoculation
of SCLC cells into immunocompromised nude mice was reduced more
efficiently in ***AAV***-***2***-infected animals compared with
tumours in mice treated with chemotherapeutic drugs alone. These data
extend and further support our previous reports on ***AAV*** functions
which might be useful in improving the efficacy of chemotherapeutic drugs
used in human cancer treatment.

L15 ANSWER 8 OF 9 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.
AN 97105758 EMBASE
DN 1997105758
TI Cellular and molecular neurosurgery: Pathways from concept to reality -
Part II: Vector systems and delivery methodologies for gene therapy of the
central nervous system.
AU Zlokovic B.V.; Apuzzo M.L.J.; Newell E.A.; Piepmeyer J.M.; Black P.M.
CS Dr. B.V. Zlokovic, 2025 Zonal Avenue, Los Angeles, CA 90033, United States
SO Neurosurgery, (1997) 40/4 (805-813).
Refs: 51
ISSN: 0148-396X CODEN: NRSRDY
CY United States
DT Journal: General Review
FS 008 Neurology and Neurosurgery
022 Human Genetics
037 Drug Literature Index
LA English
SL English
AB DIFFERENT VECTOR SYSTEMS that have been used and/or specifically developed
for central nervous system (CNS) gene transfer studies are briefly
discussed along with their advantages and disadvantages with respect to
potential clinical application. These include retroviruses, recombinant
herpes simplex virus, adenoviruses, ***adeno***-***associated***
viruses, encapsulation of plasmid deoxyribonucleic acid into
cationic liposomes, and neural and oligodendroglial stem cells.
Particular attention has been paid to relate the modality of a specific
CNS gene therapy to the strategy for adequate delivery of genetic material
to the brain for either global or localized CNS neurodegenerative chronic
disorder, as well as for CNS tumors and stroke. Techniques to circumvent
the 'impermeable' blood-brain barrier to deliver the genetic
material to the target CNS cells are reviewed and include the following:
1) local stereotactic CNS injection/infusion of viral vectors,
administration of vector producer cells, or cell replacement; 2) local
administration of genetic material into the cerebrospinal fluid
ventriculocisternal system; 3) osmotic opening of the blood-brain barrier;
4) local intra-arterial infusion; and 5) administration of blood-
brain-***tumor*** barrier permeabilizers, such as a bradykinin
B2 agonist RMP-7. It is concluded that gene therapy for several brain
disorders holds great potential, as suggested mainly by in vitro
experiments and, to some extent, by a limited number of animal
experiments. However, several drawbacks currently hamper the application
of gene therapy under the clinical setting. The problems associated with
gene therapy that still present major obstacles are as follows: 1)
inefficient transfection of host cells by viral vectors; 2) restricted
delivery of genetic material across vascular barriers of the CNS and brain
tumors; 3) nonselective expression of the transgene; and 4) in situ CNS
regulation of the transgene expression in a therapeutically controlled
manner, as imposed by the course and phenotype of the CNS disease.

L15 ANSWER 9 OF 9 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.
AN 96352402 EMBASE
DN 1996352402
TI Gene therapy against an experimental glioma using adeno-associated virus
vectors.
AU Okada H.; Miyamura K.; Itoh T.; Hagiwara M.; Wakabayashi T.; Mizuno M.;
Colosi P.; Kurtzman G.; Yoshida J.
CS Department of Neurosurgery, Nagoya University School of Medicine, 65
Tsurumai-cho, Showa-ku, Nagoya 466, Japan
SO Gene Therapy, (1996) 3/1 (957-964).
ISSN: 0969-7128 CODEN: GETHEC
CY United Kingdom
DT Journal; Article
FS 004 Microbiology
016 Cancer
022 Human Genetics
LA English
SL English
AB The efficacy of gene therapy for glioma was examined using
adeno-associated virus (***AAV***)-based vectors to deliver genes to
experimental tumors in mice. Stereotactic injection of 2 x 10⁵ U-251SP
human glioma cells into the brains of nude mice produced tumors of 19.06
± 1.78 mm² 17 days after injection. Employing a higher titer
preparation of ***AAV*** vector containing the gene for
beta-galactosidase (***AAV***-lacZ), dose-dependent transduction of
U-251SP cells was seen in vitro. When 1.6 x 10¹⁰ ***AAV***-lacZ
particles were directly injected into tumors in vivo, 30-40% of the cells
along the needle track expressed beta-galactosidase. Transduction of
U-251SP cells in vitro with an ***AAV*** vector containing a
bicistronic gene encoding both herpes simplex thymidine kinase and human
interleukin-2 (***AAV***-tk-IL-2) rendered them sensitive to the
cytotoxic effects of ganciclovir (GCV) and IL-2 was produced in a
dose-dependent manner. Cocultures of ***AAV***-tk-IL-2 transduced
cells and nontransduced cells proved highly sensitive to GCV indicating
the contribution of the bystander effect. Stereotactic delivery of 6 x
10¹⁰ ***AAV***-tk-IL-2 particles into day 7 tumors in nude mice
followed by a administration of GCV for 8 days, resulted in a 35-fold
reduction in the volume of tumors compared with controls. Normal brains
did not suffer from any toxic effect of the administration of ***AAV***
-tk-IL-2 and GCV. These results indicate that high titer ***AAV***
vector treatment may be safe and effective for in vivo gene therapy of
human brain tumors.

=> d his
(FILE 'HOME' ENTERED AT 16:29:22 ON 02 JAN 2002)
FILE 'BIOSIS, EMBASE, CAPLUS' ENTERED AT 16:29:48 ON 02 JAN 2002
L1 3153 S ADENO-ASSOCIATED VIRUSES OR AAV
L2 3153 S ADENO ASSOCIATED VIRUSE? OR AAV
L3 536 S ADENO ASSOCIATED VIRUS 2 OR AAV-2
L4 3233 S L1 OR L3
L5 201 S L4 AND CANCER?
L6 5 S L4 AND (CHEMO)? RESISTAN?
L7 2 DUP REM L6 (3 DUPLICATES REMOVED)
FILE 'STNGUIDE' ENTERED AT 16:38:04 ON 02 JAN 2002
FILE 'EMBASE, BIOSIS, CAPLUS' ENTERED AT 16:38:10 ON 02 JAN 2002
L8 16 S L4 AND (COLON CANCER OR PANCREATIC CARCINOMA OR BRAIN TUMOR
L9 0 S L4 AND RADIOTHERAPY AND RESISTANCE
L10 170 S L4 AND RESISTANCE
L11 1 S L10 AND (CHEMO OR RADIOT?)
L12 5 S L10 AND RADIATION
L13 6 S L11 OR L12
L14 4 DUP REM L13 (2 DUPLICATES REMOVED)
L15 9 DUP REM L8 (7 DUPLICATES REMOVED)
=> s l4 and (cisplatin or etoposide)
L16 29 L4 AND (CISPLATIN OR ETOPOSIDE)
=> dup rem l16
PROCESSING COMPLETED FOR L16
L17 14 DUP REM L16 (15 DUPLICATES REMOVED)
=> d bib abs 1-
YOU HAVE REQUESTED DATA FROM 14 ANSWERS - CONTINUE? Y(N):y
L17 ANSWER 1 OF 14 CAPLUS COPYRIGHT 2002 ACS
AN 2001:798037 CAPLUS
DN 135:339221
TI Cytotoxic single-stranded DNA agents
IN Raj, Kenneth; Beard, Peter Martin
PA BTG International Limited, UK
SO PCT Int. Appl., 51 pp.
CODEN: PIXXD2
DT Patent
LA English
FAN.CNT 1
PATENT NO. KIND DATE APPLICATION NO. DATE
PI WO 2001080840 A2 20011101 WO 2001-GB1795 20010420
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM,
HR, HU, ID, IL, IN, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,
LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO,
RU, SD, SE, SG, SI, SK, SL, TJ, TM, TT, TZ, UA, UG, US, UZ,
VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,
DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,
BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG
PRAI GB 2000-0887 A 20000420
AB A method of killing a cell that is lacking in effective p53 protein
activity, particularly as compared to wild type, is provided. It is
characterized in that it comprises delivering to the cell a single
stranded DNA including a portion with at least one base, internally
located with respect to any 3' and 5' ends of the DNA, that is
unbasepaired with another base in a form that is capable of being
internalised by the cell.

L17 ANSWER 2 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 1

AN 2001325528 EMBASE

TI Adeno-associated virus type 2-mediated transduction of human monocyte-derived dendritic cells: Implications for ex vivo immunotherapy.

AU Ponnazhagan S.; Mahendra G.; Curiel D.T.; Shaw D.R.

CS S. Ponnazhagan, Department of Pathology, LHRB 513, University of Alabama, 701 18th St. South, Birmingham, AL 35294-0007, United States.

sponnazh@path.uab.edu
SO Journal of Virology, (2001) 75/19 (9493-9501).

Refs: 56

ISSN: 0022-538X CODEN: JOVIAM

CY United States

DT Journal Article

FS 004 Microbiology

026 Immunology, Serology and Transplantation

LA English

SL English

AB Dendritic cells (DCs) are pivotal antigen-presenting cells for regulating immune responses. A major focus of contemporary vaccine research is the genetic modification of DCs to express antigens or immunomodulatory molecules, utilizing a variety of viral and nonviral vectors, to induce antigen-specific immune responses that ameliorate disease states as diverse as malignancy, infection, autoimmunity, and allergy. The present study has evaluated adeno-associated virus (AAV) type 2 as a vector for ex vivo gene transfer to human peripheral blood monocyte (MO)-derived DCs. AAV is a nonpathogenic parvovirus that infects a wide variety of human cell lineages in vivo and in vitro, for long-term transgene expression without requirements for cell proliferation. The presented data demonstrate that recombinant AAV (rAAV) can efficiently transduce MOs as well as DCs generated by MO culture with granulocyte-macrophage colony-stimulating factor plus interleukin in vitro. rAAV transgene expression in MO-derived DCs could be enhanced by AAV-epitope, previously reported to enhance AAV gene expression. rAAV transduction of freshly purified MO followed by 7 days of culture with cytokines to generate DCs, and subsequent sorting for coexpression of DC markers CD1a and CD40, showed robust transgene expression as well as evidence of nuclear localization of the rAAV genome in the DC population. Phenotypic analyses using multiple markers and functional assays of one-way allogeneic mixed leukocyte reactions indicated that rAAV-transduced MO-derived DCs were as equivalent to nontransduced DCs. These results support the utility of rAAV vectors for future human DC vaccine studies.

L17 ANSWER 3 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 2

AN 2000026402 EMBASE

TI Transduction of hepatocellular carcinoma (HCC) using recombinant adeno-associated virus (AAV): In vitro and in vivo effects of genotoxic agents.

AU Peng D.; Qian C.; Sun Y.; Barajas M.A.; Prieto J.

CS C. Qian, Div. of Hepatology and Gene Therapy, Department of Internal Medicine, University of Navarra, 31080 Pamplona, Spain. cqian@unav.es
SO Journal of Hepatology, (2000) 32/6 (975-985).

Refs: 51

ISSN: 0168-8278 CODEN: JOHEEC

CY Denmark

DT Journal Article

FS 016 Cancer

022 Human Genetics

037 Drug Literature Index

048 Gastroenterology

LA English

SL English

AB Background/Aims: Adeno-associated virus (AAV) is an attractive tool for gene therapy. Here we investigated the in vitro and in vivo transduction of hepatocellular carcinoma (HCC) cells by an AAV vector and the efficacy of different strategies to enhance the transduction of the tumor. Methods: Transduction efficiency was determined by analyzing AAV-mediated beta-galactosidase gene (AAVlacZ) expression. Results: Adenovirus help or pretreatment of HCC cells with gamma-irradiation or with the topoisomerase inhibitor etoposide resulted in marked enhancement of cell transduction in vitro. In vivo studies in nude mice with subcutaneous HCC tumors showed that HCC cells were not transduced by AAV vector alone. However, co-infection of the tumor with adenovirus allowed an efficient expression of the reporter gene but only at the sites of vector injection. Previous gamma-irradiation of subcutaneous tumors with 1800 rad was able to improve transduction of HCC cells (up to 30%) using recombinant AAV. Continuous i.p. infusion of etoposide in buffalo rats harboring HCC tumors in the liver resulted in transduction of normal liver tissue and also of very small neoplastic lesions (<2 mm) but no transduction was observed in tumors bigger than 2 mm. To analyze this phenomenon we determined etoposide concentration in hepatic tissue. Our results revealed high concentrations of the drug in non-tumoral tissue but almost undetectable levels in big tumor nodules. Conclusions: Our results indicate that while both radiotherapy and etoposide enhance transduction of tumor cells by AAV in vitro, only radiotherapy increases tumor transduction in vivo. Our data suggest the existence of a barrier which limits in vivo the diffusion of chemotherapeutic agents to well-established HCC nodules.

L17 ANSWER 4 OF 14 CAPLUS COPYRIGHT 2002 ACS

AN 1999764174 CAPLUS

DN 132.9629

TI Adeno-associated viral vector-mediated expression of factor VIII activity

IN Cohen, Lawrence K.; Spratt, S. Kaye; Couto, Linda

PA Cell GeneSys, Inc., USA

SO PCT Int. Appl., 38 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

PATENT NO. KIND DATE APPLICATION NO. DATE

PI WO 9961595 A2 19991202 WO 1999-US10472 19990527

WO 9961595 A3 20000127

W: AE, AL, AM, AT, AU, AZ, BA, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG,

CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG

AU 9941856 A1 19991213 AU 1999-41856 19990527

EP 1082445 A2 20010314 EP 1899-025606 19990527

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI

PRAI US 1999-84423 A 19990527

WO 1999-US10472 W 19990527

AB The invention demonstrates that recombinant AAV (rAAV) vectors may be used to deliver for effective expression a protein with Factor VIII function to treat hemophilia A. The invention provides methods and materials for expressing polypeptides with factor VIII activity comprising administering at least two rAAV vectors encoding different domains of human factor VIII and at least the heavy and light chains.

L17 ANSWER 5 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.

AN 1999113632 EMBASE

TI Enhancement of UV-induced cytotoxicity by the adeno-associated virus replication proteins.

AU Zhou C.; Yang Q.; Trempe J.P.

CS J.P. Trempe, Dept. of Biochemistry/Molec. Biol., Medical College of Ohio, 3035 Arlington Ave., Toledo, OH 43614-5804, United States. jtrempe@mco.edu

SO Biochimica et Biophysica Acta - Gene Structure and Expression, (19 Mar 1999) 1444/3 (371-383).

Refs: 80

ISSN: 0167-4781 CODEN: BBSGDS

PUI S 0167-4781(99)00014-7

CY Netherlands

DT Journal Article

FS 029 Clinical Biochemistry

LA English

SL English

AB Adeno-associated virus (AAV) normally requires co-infection of a helper virus to complete its life cycle. However, under conditions of cellular stress, such as treatment with carcinogens or ultraviolet (UV) light, a permissive intracellular environment is established and AAV completes its replicative cycle producing low levels of progeny virus. AAV DNA replication is dependent upon viral replication proteins, Rep78 and Rep68. The detailed mechanism by which these proteins interact with host cell factors is unknown. We have used a cell line (Neo6) that inducibly expresses the AAV Rep proteins to study their effects on cells that have undergone UV-induced DNA damage. Induction of Rep protein expression immediately after a sub-lethal dose of UV irradiation resulted in rapid cell killing. Those cells that die had chromatin condensation while cellular membranes remained intact, suggesting that concurrent Rep expression and UV damage induces an apoptosis-like response. However, we did not observe any DNA degradation. Thus we believe that the combination of Rep expression and UV irradiation induces cell death that shares some of the characteristics of apoptosis. UV irradiation and Rep expression induced an increase in the level of the CDK inhibitor, p21(Cip), and the appearance of modified forms of both p21(Cip) and Bcl-2. Alteration of normal expression of these cytoskeletal/apoptotic proteins provides insight into the intracellular targets of the AAV replication proteins.

L17 ANSWER 6 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 3

AN 1999062500 EMBASE

TI Enhanced sensitivity of small cell lung cancer cell lines to cisplatin and etoposide after infection with adeno-associated virus type 2.

AU Hüggenberg M.; Schlehofer J.R.; Von Knebel Doeberitz M.; Klein-Baernschmitt P.

CS P. Klein-Baernschmitt, Sek. Molekuläre Diagnostik Therapie, Chirurgische Universitätsklinik, Universität Heidelberg, 69120 Heidelberg, Germany.

p.klein@dkfz-heidelberg.de

SO European Journal of Cancer, (1999) 35/1 (108-110).

Refs: 21

ISSN: 0959-8049 CODEN: EJCAEL

PUI S 0959-8049(99)00275-5

CY United Kingdom

DT Journal Article

FS 004 Microbiology

015 Chest Diseases, Thoracic Surgery and Tuberculosis

016 Cancer

037 Drug Literature Index

LA English

SL English

AB In previous studies we have reported the sensitisation of human tumour cells to gamma irradiation and chemotherapeutic drugs upon infection with the human non-pathogenic adeno-associated virus type 2 (AAV-2) in vitro and in vivo. Treatment of small cell lung cancer (SCLC) is consistently hampered by relapses due to the selection of chemotherapy-resistant cell clones. Hence, we were interested to test whether selection of chemotherapy-resistant SCLC cells might be reduced or even prevented if chemotherapy is applied in combination with AAV-2 infection. In vitro proliferation assays indicated that the number of proliferating cells, after combined treatment with cisplatin and etoposide, can be significantly reduced by concomitant AAV-2 infection, as compared with treated but non-infected controls. H460 SCLC cells, which show resistance to etoposide / cisplatin chemotherapy (compared with a cell line which was never chemotherapeutically treated before, like NCH-H209) were significantly more sensitive after AAV-2 infection. Infection, suggesting that the therapeutic efficacy of chemotherapy in SCLC can be enhanced even if the cells are already relatively resistant to chemotherapy. Similarly, in vivo growth of tumors induced by inoculation of SCLC cells into immunocompromised nude mice was reduced more efficiently in AAV-2 infected animals compared with tumours in mice treated with chemotherapeutic drugs alone. These data extend and further support our previous reports on AAV functions which might be useful in improving the efficacy of chemotherapeutic drugs used in human cancer treatment.

L17 ANSWER 7 OF 14 CAPLUS COPYRIGHT 2002 ACS

AN 1998734970 CAPLUS

DN 129-340538

TI Increasing transduction of cells by adeno-associated virus vectors by using DNA metabolism-altering agents

IN Alexander, Ian E.; Russell, David W.; Miller, A. Dusty

PA Fred Hutchinson Cancer Research Center, USA

SO U.S., 12 pp. Cont.-in-part of U.S. 5,804,080.

CODEN: USXXAM

DT Patent

LA English

FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PI	US	5834182	A	19981110	US	1987-750274	19970225
	US	5804080	A	19870218	US	1994-254312	19940606
	WO	9533824	A1	19951214	WO	1995-US7202	19950605

W: AU, CA, JP, US

RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE
PRAIUS 1994-254312 19940606
WO 1995-US7202 19950605

AB This invention includes methods for increasing the efficiency of transduction of cells, including non-dividing cells, by recombinant adeno-associated virus ("AAV") vectors. The methods utilize agents that alter certain aspects of DNA metab., more specifically, that affect DNA synthesis and/or affect repair, that impact on maintenance of chromosomal integrity, and/or that cause damage to the cellular DNA. Agents and vectors can now also be preselected and screened for transducing ability and/or transducing agents for their effect on DNA metab. These agents include irradiated nucleotides such as thymidine, gamma irradi., UV irradi., cis-platinum, "etoposide", hydroxyurea and aphidicolin.

L17 ANSWER 8 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.
AN 1998180472 EMBASE

TI Adeno-associated virus gene transfer into renal cells: Potential for in vivo gene delivery

AU Langer J.C.; Klotman M.E.; Hanss B.; Tulin N.; Bruggeman L.A.; Klotman P.E.; Upkowitz M.S.

CS Dr. M.S. Upkowitz, Nephrology Division, Box 1243, Mount Sinai School of Medicine, 1 Gustave L. Levy Place, New York, NY 10029, United States.
mlupkow@msm.mssm.edu

SO Experimental Nephrology, (1998) 6/3 (189-194).

Refs: 24

ISSN: 1018-7782 CODEN: EXNEEG

CY Switzerland

DT Journal: (Short Survey)

FS 004 Microbiology

028 Urology and Nephrology

029 Clinical Biochemistry

LA English

SL English

AB The human parvovirus adeno-associated virus ("AAV"), type 2, has a number of features that make it an attractive choice as a vector for gene delivery to the kidney. "AAV" vectors permit long-term gene expression in vivo by integration into the host genome, have potential for site-specific integration of chromosome 19, do not express viral genes or generate a cellular immune response, and demonstrate enhancement of gene expression by chemotherapeutic agents that are approved for use in vivo. These properties confer advantages to "AAV" over other viral and nonviral methods for gene transfer. Preliminary experiments in our laboratory suggest that "AAV" is able to transfer genes to both renal cells in culture and the kidney in vivo. Thus, "AAV" has the potential to be an important gene transfer vector for the kidney in vivo.

L17 ANSWER 9 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 4

AN 97217501 EMBASE

DN 1997217501

TI Transduction by adeno-associated virus vectors in the rabbit airway: Efficiency, persistence, and readministration

AU Halbert C.L.; Standert T.A.; Alken M.L.; Alexander I.E.; Russell D.W.; Miller A.D.

CS A.D. Miller, Fred Hutchinson Cancer Research Ctr., 1100 Fairview Ave. North, Seattle, WA 98108, United States. dmiller@fhrc.org
SO Journal of Virology, (1997) 71/8 (5932-5941).

Refs: 44

ISSN: 0022-538X CODEN: JOVIAH

CY United States

DT Journal: Article

FS 004 Microbiology

022 Human Genetics

LA English

SL English

AB The ability of recombinant adeno-associated virus ("AAV") vectors to integrate into the host genome and to transduce nondividing cells makes them attractive as vehicles for gene delivery. In this study, we assessed the ability of several "AAV" vectors to transduce airway cells in rabbits by measuring marker gene expression. "AAV" vectors that transferred either a beta-galactosidase (beta-gal) or a human placental alkaline phosphatase (AP) gene were delivered to one lobe of the rabbit lung by use of a balloon catheter placed under fluoroscopic guidance. We observed vector-encoded beta-gal or AP staining almost exclusively in the epithelial and smooth muscle cells in the bronchus at the region of balloon placement. The overall efficiency of transduction in the balloon-treated bronchial epithelium was low but reached 20% in some areas. The majority of the staining was in ciliated cells but was also observed in basal cells and airway smooth muscle cells. We observed an 80-fold decrease in marker-positive epithelial cells during the 60-day period after vector infusion, whereas the number of marker-positive smooth muscle cells stayed constant. Although treatment with the topoisomerase inhibitor "etoposide" dramatically enhanced "AAV" transduction in primary airway epithelial cells in culture, treatment of rabbits did not improve transduction rates in the airway. Vector readministration failed to produce additional transduction events, which correlated with the appearance of neutralizing antibodies. These results indicate that both readministration and immune modulation will be required in the use of "AAV" vectors for gene therapy to the airway epithelium.

L17 ANSWER 10 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 5

AN 97081784 EMBASE

DN 1997081784

TI Persistent expression of human clotting factor IX from mouse liver after intravenous injection of adeno-associated virus vectors

AU Koeberl D.D.; Alexander I.E.; Halbert C.L.; Russell D.W.; Miller A.D.
CS A.D. Miller, Fred Hutchinson Can. Res. Center, 1100 Fairview Avenue North, Seattle, WA 98108, United States. dmiller@fhrc.org

SO Proceedings of the National Academy of Sciences of the United States of America, (1997) 94/4 (1428-1431).

Refs: 43

ISSN: 0027-8424 CODEN: PNASAB

CY United States

DT Journal: Article

FS 022 Human Genetics

025 Hematology

LA English

SL English

AB We previously found that gene transduction by adeno-associated virus ("AAV") vectors in cell culture can be stimulated over 100-fold by treatment of the target cells with agents that affect DNA metabolism, such as irradiation or topoisomerase inhibitors. Here we show that previous gamma-irradiation increased the transduction rate in mouse liver by up to 900-fold, and the topoisomerase inhibitor "etoposide" increased transduction by about 20-fold. Similar rates of hepatic transduction were obtained by direct injection of the liver or by systemic delivery via tail vein injection. Hepatocytes were much more efficiently transduced than other cells after systemic delivery, and up to 3% of all hepatocytes could be transduced after one vector injection. The presence of wild-type "AAV" , which contaminates many "AAV" vector preparations, was required to observe a full response to gamma-irradiation. Injection of mice with "AAV" vectors encoding human clotting factor IX after gamma-irradiation resulted in synthesis of low levels of human clotting factor IX for the 5-month period of observation. These studies show the potential of targeted gene transduction of the liver by "AAV" vectors for treatment of various hematological or metabolic diseases.

L17 ANSWER 11 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 6

AN 96284696 EMBASE

DN 1996284696

TI Improved efficacy of chemotherapy by parvovirus-mediated sensitisation of human tumour cells

AU Klein-Baumenschmidt P.; Von Knebel Doeberitz M.; Ehrbar M.; Geletnek K.; Kleinschmidt J.; Schlehofer J.R.

CS German Cancer Research Center, Applied Tumour Virology, 69120 Heidelberg, Germany

SO European Journal of Cancer Part A, (1996) 32/10 (1774-1780).

ISSN: 0959-8049 CODEN: EJCTEA

CY United Kingdom

DT Journal: Article

FS 004 Microbiology

018 Cancer

030 Pharmacology

037 Drug Literature Index

LA English

SL English

AB Increasing resistance of tumour cells towards the cytotoxic action of chemotherapeutic drugs is a major limitation in the treatment of cancer patients. The non-pathogenic human "adeno" - "associated" "viruses" ("AAV") have been reported to sensitise HeLa cervical cancer cells to gamma irradiation in vivo and in vitro. To test whether these parvoviruses might render other human tumour cells more sensitive towards chemotherapeutic drugs, we analysed the effects of "AAV" type 2 ("AAV" - "2") infection on established cancer cell lines and freshly explanted tumour biopsies treated with chemotherapeutic agents (e.g. "cisplatin") - "AAV" - "2" infection significantly increased the cytotoxic activity of chemotherapeutic drugs compared with uninfected controls. "AAV" - "2" infection without concomitant chemotherapeutic treatment had no significant effect on viability of the cells. In nude mice, combined application of "AAV" - "2" infection and chemotherapeutic treatment significantly increased the therapeutic activity on tumours arising from subcutaneously injected tumour cells compared with tumours treated by chemotherapeutics only. These results indicate that "AAV" - "2" infection sensitises human cancer cells towards the cytotoxic action of chemotherapeutic drugs.

L17 ANSWER 12 OF 14 CAPLUS COPYRIGHT 2002 ACS

AN 199687115 CAPLUS

DN 124108980

TI Increasing transduction of cells by adeno-associated virus vectors by using DNA metabolism-altering agents

IN Alexander, Ian E.; Russell, David W.; Miller, A. Dusty.

PA Fred Hutchinson Cancer Research Center, USA

SO PCT Int. Appl. 29 pp.

CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 2

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

PI	WO	9533824	A1	19951214	WO	1995-US7202	19950605
	W:	AU, CA, JP, US					
	RW:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE					
	US	5804080	A	19870218	US	1994-254312	19940606
	CA	2192214	AA	19951214	CA	1995-2192214	19950605
	AU	9526994	A1	19960104	AU	1995-26994	19950605
	EP	765387	A1	19970402	EP	1995-922237	19950605
	R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE					
	PRAIUS	1994-254312	A	19871110	US	1987-750274	19970225
	WO	1995-US7202					

AB Methods are provided for increasing the efficiency of transduction of cells, including non-dividing cells, by recombinant "AAV" vectors. The methods utilize agents that alter certain aspects of DNA metab., more specifically, that affect DNA synthesis and/or affect repair, that impact on maintenance of chromosomal integrity, and/or that cause damage to the cellular DNA. Agents and vectors can now also be preselected and screened for transducing ability and/or transducing agents for their effect on DNA metab. These agents include irradiated nucleotides such as thymidine, gamma irradi., UV irradi., cis-platinum, "etoposide", hydroxyurea, aphidicolin, and camptothecin.

L17 ANSWER 13 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 7

AN 95185325 EMBASE

DN 1995185325

TI DNA synthesis and topoisomerase inhibitors increase transduction by adeno-associated virus vectors

AU Russell D.W.; Alexander I.E.; Miller A.D.

CS Fred Hutchinson Cancer Research Ctr., Seattle, WA 98104, United States

SO Proceedings of the National Academy of Sciences of the United States of America, (1995) 92/12 (5719-5723).

ISSN: 0027-8424 CODEN: PNASAB

CY United States

DT Journal: Article

FS 021 Developmental Biology and Teratology

LA English

SL English

AB Viral vectors based on adeno-associated virus (***AAV***)

preferentially transduce cells in S phase of the cell cycle. We recently found that DNA-damaging agents increased the transduction of nondividing cells. However, the optimal concentrations were toxic to cells. Here we show that the transduction of normal human fibroblasts by ***AAV*** vectors is increased by prior exposure to DNA synthesis inhibitors, such as aphidicolin or hydroxyurea, and topoisomerase inhibitors, such as "etoposide" or camptothecin. Transduction efficiencies could be increased >300-fold in stationary cultures at concentrations that did not affect cell viability or proliferative potential. Both S-phase and non-S-phase cells were affected, suggesting that cellular functions other than replicative DNA synthesis may be involved. Applying these methods to gene transfer protocols should improve prospects for gene therapy by ***AAV*** vectors.

L17 ANSWER 14 OF 14 EMBASE COPYRIGHT 2002 ELSEVIER SCI. B.V.DUPLICATE 8

AN 90340734 EMBASE

DN 1990340734

TI Induction of selective DNA amplification and morphological cell transformation in Syrian hamster embryo cells.

AU Klein P.; Poot-Zobel B.L.; Yalkinoglu O.; Schlehofer J.R.

CS German Cancer Research Center, Inst. of Chemotherapy/Toxicol., Im Neuenheimer Feld 280, D-6900 Heidelberg, Germany

SO Mutation Research, (1990) 232/2 (183-190).

CODEN: MUREAV

CY Netherlands

DT Journal: Article

FS 018 Cancer

022 Human Genetics

052 Toxicology

037 Drug Literature Index

LA English

SL English

AB DNA amplification is a frequently observed event in continuous cell lines and in tumors. It is likely that a common mechanism underlies the amplification of specific DNA sequences which confer drug resistance and genes which give a growth advantage to the tumor. To find a correlation between the induction of DNA amplification by chemicals and morphological cell transformation we treated Syrian hamster embryo (SHE) cells with diverse antineoplastic agents of different classes. Analysis of these agents seems to be important since they are potentially carcinogenic and resistance inducing. For the measurement of DNA amplification we established a new system using adeno-associated virus type 2 (***AAV***)-infected primary SHE cells as target cells and amplification of viral DNA as marker of DNA amplification. Simultaneously we determined morphological cell transformation in SHE cells. Our findings demonstrate that there is only a limited correlation between the induction of ***AAV*** DNA amplification and the morphological cell transformation in SHE cells. The newly established system of ***AAV*** DNA amplification appears to be a useful tool for the investigation of drug resistance in target cells of choice.

=>

--Logging off of STN--

=>

Executing the logoff script...

=> LOG Y

COST IN U.S. DOLLARS	ENTRY	SINCE FILE	SESSION	TOTAL
FULL ESTIMATED COST		101.00		142.72
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	ENTRY	SINCE FILE	SESSION	TOTAL
CA SUBSCRIBER PRICE		-4.34		-4.34

STN INTERNATIONAL LOGOFF AT 16:48:42 ON 02 JAN 2002